

CAVOR GEL

Gel ready to fight c GAINST cockroaches and ants

Contains 0.05% Fipronil

Product presentation

Trade name:

CAVOR GEL

Active ingredient:

Fipronil 0.05%

Description and Mode of Action:

CAVOR GEL is effective against German cockroaches (German cockroaches), oriental cockroaches (oriental cockroaches) and American cockroaches (*periplaneta Americana*) and black or garden ants (*Lasius niger*), yellow ants (*Lasius lavus*), *Lasius* ants red (*Lasius emarginatus*) against both nymphs and adults.

CAVOR GEL can be used against cockroaches and ants in homes (including kitchens). Public buildings such as hospitals, prisons, theaters, hotels, etc. Industrial and commercial premises such as factories, stores, restaurants, and commercial kitchens, workshops, aircraft cabins, vehicles, land and boats

Manual :

CAVOR GEL is most effective when placed in or near places infested by cockroaches and ants , in places where they search or deposit food. These places are usually dark, hot, humid and quiet, for example behind refrigerators and in electrical installations. He false to seek these places by a systematic inspection or better by trapping. Better results are obtained if the surface to be treated is stored beforehand before treatment and if the precaution is taken to remove all other sources of food. CAVOR GEL should be applied to infested areas in the form of small drops. Gel Drops should be applied in dark, hard-to-reach crevices, cracks, behind cupboards and shelves, under and in kitchen appliances, and in electricity meters, garbage chutes, pipes and bathroom installations , etc. Do not apply CAVOR GEL in a place where it could be flooded or eliminated by the household. Do not spray other insecticides on or around CAVOR GEL, nor place it on areas already treated, which could reduce the attractiveness of the gel.

If CAVOR GEL spills onto surfaces not intended to be treated, it should be removed with a damp paper towel and disposed of in the garbage. Clean the surface thoroughly.

Cockroaches and ants die a few hours after a single intake of CAVOR GEL. Dead cockroaches and ants will be found in infested places 24 hours after treatment.

Treated surfaces will need to be re-inspected after 1 to 2 weeks. In the event of heavy infestation, a second application of CAVOR GEL may be necessary if the first treatment has been consumed and cockroaches and ants remain.

Application doses:

In case of heavy infestation and in particularly dirty and / or congested places or if it is not possible to remove all other food sources, a higher dose should be used.

Normal dose:

German cockroach: one drop of 0.03 g gel per square meter

Oriental or American cockroach: two drops of 0.03g gel per square meter.

Ants : One drop of 0.03 g (3 - 4 mm in diameter) every 30 cm on the path.

High dose:

German cockroach: two drops of 0.03 g gel per square meter

Oriental or American cockroach: three drops of 0.03 g gel per square meter

Ants : One drop of 0.03 g (3 - 4 mm in diameter) every 15 cm on the path.

Storage:

Store CAVOR GEL cartridges in the original box away from large variations in temperature.

Precautionary advice:

S2 - keep out of reach of children

S13 - keep away from food and drink, including animal feed

S20 / S21 - Do not eat, drink or smoke while using.

Dispose of empty packaging only by taking all the usual precautions

Do not try to open the cartridges or refill them.

Do not put gel in places where food and water could be contaminated.

Use only in locations inaccessible to children and animals

Do not use outdoors.